WORKING PAPERS SERIES
DIPARTIMENTO DI
SCIENZE SOCIALI ED ECONOMICHE
n. 6/2015

**Estimating Non-Linear DSGEs with the
Approximate Bayesian Computation:
an application to the Zero Lower Bound**

Author:

Valerio Scalone

Estimating Non-Linear DSGEs with the Approximate Bayesian Computation: an application to the Zero Lower Bound

Valerio Scalone*

November 8, 2015

Abstract

Non-linear model estimation is generally perceived as impractical and computationally burdensome. This perception limited the diffusion on non-linear models estimation. In this paper a simple set of techniques going under the name of Approximate Bayesian Computation (ABC) is proposed. ABC is a set of Bayesian techniques based on moments matching: moments are obtained simulating the model conditional on draws from the prior distribution. An accept-reject criterion is applied on the simulations and an approximate posterior distribution is obtained by the accepted draws. A series of techniques are presented (ABC-regression, ABC-MCMC, ABC-SMC). To assess their small sample performance, Montecarlo experiments are run on AR(1) processes and on a RBC model showing that ABC estimators outperform the Limited Information Method (Kim, 2002), a GMM-style estimator. In the remainder, the estimation of a new-keynesian model with a zero lower bound on the interest rate is performed. Non-gaussian moments are exploited in the estimation procedure.

*University of Rome, La Sapienza, Department of Social Sciences and Economics, LUISS Guido Carli, Department of Economics. Email: vsalone@luiss.it. I thank Salvatore Nistico' and Giuseppe Ragusa for their support. I also thank Pierpaolo Benigno, Nicola Borri, Nuno Coimbra, Sergio de Ferra, Giorgio Di Giorgio, Jesus Fernandez-Villaverde, Maddalena Galardo, Raffella Giacomini, Anastasios Karantounias, Lena Koerber, Dennis Kristensen, Riccardo Masolo, Marco Lippi, Lucrezia Reichlin, Pietro Reichlin, Federica Romei, Frank Schorfheide, Paolo Surico for useful comments. I am also grateful to seminar participants at LUISS Guido Carli, the London Business School, UCL, University of Pennsylvania.

JEL CLASSIFICATION: C15, C11, E2

Key words: Monte-Carlo analysis; Method of moments, Bayesian, Zero Lower Bound, DSGE estimation

1 Introduction

DSGE (Dynamic stochastic general equilibrium) models play an important role in Macroeconomic theory. In the last decade, they became the workhorse of many central banks. They are used to explain economic fluctuations from a general equilibrium perspective, to make forecasts on the path of macroeconomic variables, to advise policy makers in taking decisions. Model estimation is a crucial step allowing economists to make quantitative statements in the framework of a probabilistic structure.

Great moderation years have seen the prevalence of linear methods: linearization to solve the model, Kalman filter to compute the likelihood and Bayesian techniques to estimate the model. The Great Recession, the presence of a lower bound reached by the policy interest rate, the general increase in volatility, the need to model a fraction of borrowing constrained households pushed researchers to inquire about the role of non-linearities in the economic models. Log-linearization and Kalman filter are not fit to represent some features of the data (presence of occasionally binding constraints, stochastic volatility, non-Gaussian shocks) and non-linear solution methods are being developed. The Particle filter is the method usually applied in the estimation of non-linear models (Fernandez-Villaverde et al.). The Particle filter is computationally burdensome, especially to handle medium or large-scale DSGE models. Besides, the Particle filter needs the introduction measurement errors, to avoid the degeneracy of the particles and compute the likelihood. In many cases, given the size of the model, the standard deviation of the measurement errors is fixed in advance. All these issues limited the diffusion of non-linear estimation so far.

In this paper, Approximate Bayesian Computation (ABC), a set of techniques based on simulation and moments matching, is proposed as an alternative to estimate non-linear models. ABC techniques are presented. Two Montecarlo experiments on ABC methods and the Bayesian Limited Information Method (BLI) are assessed. The goal is comparing the small sample performance of the two estimators. Moreover, ABC is applied to the estimation of standard new-keynesian model with an occasional binding positivity constraint on the interest rate.

Approximate Bayesian Computation techniques are a set of techniques developed

in natural sciences. The core mechanism in ABC (ABC-rejection) is the following:

- The model is simulated a large number of times, conditional on the vectors of parameters drawn from the prior distribution. Each simulation has the same sample size of the observed sample;
- Euclidean distance between the moments of each simulation and the observed ones is computed for each simulation;
- Each simulation is accepted or rejected if the Euclidean distance is below or above a tolerance level;
- The accepted draws are a sample of the approximate posterior distribution.

Drawing from the prior distribution can be very inefficient if the prior and the posterior distributions are very different. This causes very low acceptance ratios and may make simple ABC-rejection impractical.

To tackle the computational inefficiency, a series of refinements have been developed:

- ABC-regression: the accepted draws are corrected with a post-sampling correction step;
- ABC-MCMC: the accept-reject is applied to explore the posterior distribution building a Markov Chain;
- ABC-SMC: the draws are iteratively sampled from the approximate posterior distribution.

In Economics, the estimator proposed by Creel and Kristensen (2013) in its Bayesian simulated version (Simulated Bayesian Indirect Likelihood estimator, SBIL) coincides with a variant of ABC (ABC-kernel). Creel and Kristensen provide asymptotic results for the estimator, compare the small sample performance of the estimator with the Simulated Method of Moments from a frequentist perspective: they compute the RMSEs with respect to the true values. They also apply the method in the estimation of a baseline DSGE model, solved with perturbation methods.

Instead, in this paper, the comparison is done between ABC methods and the Bayesian Limited Information Method (BLI). The BLI is a Bayesian method based on exploiting the likelihood of the moments. It can be intuitively thought as the Bayesian version of the Generalized Method of Moments and the Simulated Method of Moments. In DSGE estimation, it has been applied by Christiano, Trabandt and

Walentin,(2010), and Christiano, Eichenbaum and Trabandt (2014) and it is getting more and more popular among researchers.

The comparison of small sample performance is done from a Bayesian perspective: the RMSEs are computed with respect to the Full likelihood posterior mean and the approximate posterior distribution are compared to the Full likelihood posterior distributions. The Montecarlo experiments are run using an AR(1) model and a RBC model with three observables and identification issues. The persistence and the sample sizes of the models are diverse to check the different performances of the estimators.

ABC estimators outperform BLI estimator using small samples and high persistence processes. With large samples, they have the same performance, provided that the number of simulations is sufficiently large to get rid of the simulation effect. This hold both for the AR(1) and the RBC model.

BLI and GMM-style estimators exploit the information contained in the moments. GMM-style estimators build the likelihood function/objective function relying on the normality assumption of the moments distribution: moments and their variances are sufficient statistics of asymptotically normally distributed moments. Instead, ABC estimators explore the whole distribution of the moments. This is a comparative advantage with respect to the BLI estimator, especially when the distribution of the moments is far from being normal and not centred around the population moment. This difference is more remarkable with small samples and high persistence. In that case, convergence of moments to the normal distribution is slower and the actual moments distribution substantially differs from the asymptotic distribution.

For this same type of reason, ABC can exploit non-Gaussian moments: binomially and multinomially distributed moment. As an example, in an estimation procedure of an economic models, ABC techniques can try to match the frequency of recessions (and expansion), of deflation (and inflation) and so forth.

These results paved the way for a real life application: the estimation of a newkeynesian model with occasionally binding positivity constraint (models with Zero Lower Bound, ZLB). Models with occasionally binding constraints produce moments which do not respect the regularity assumption requested to apply GMM-style estimators. ABC can estimate such models, since the moments distribution is explored through the accept-reject method, taking into account the actual distribution of the moments. Moreover, ABC permits to match non-gaussian moments: the probability of being at the ZLB, the number of episodes and so forth. The non-linearity generated by the occasionally binding constraint and the gap between the

notional interest rate and the zero lower bound is handled by fully non-linear methods or piecewise linear methods. Perturbation methods (log-linearization, 2nd order approximation and so forth) cannot handle the solution of a model with occasionally binding constraint, since they approximate the solution around a steady state in which the zero lower bound is not binding.

In this paper, a piecewise linear approximation method is applied to the solution of a model with ZLB. The model is estimated according to an ABC-Sequential Montecarlo technique. ABC-SMC is helpful to tackle the curse of dimensionality increasing the acceptance ratio. The estimates are exploited to produce some consideration about the role of the ZLB in the economy.

Summing up, the contributions of this work are the following. ABC techniques are explored and applied to the estimation of economic models. Moreover, a comparison with the Bayesian Limited Information is assessed from a Bayesian perspective: ABC estimators outperform GMM-style estimators in terms of RMSE (computed with respect to the Full likelihood estimator). This is particularly true dealing with small samples and highly persistent processes. Besides, the estimation of a model with a Zero Lower Bound is performed, using gaussian and non-gaussian moments. The estimation is performed using six observable variables and a dataset including 2013Q3¹.

The reminder of the paper is the following. Section 2 presents the ABC techniques. In Section 3 a comparison between ABC estimator and the BLI estimator is assessed. In Section 4 the model with ZLB is estimated. Section 5 concludes.

2 Approximate Bayesian Computation.

The Approximate Bayesian Computation (ABC) is a set of statistical techniques developed in population genetics at the end of the 90's (Pritchard, 2000). In the last decade, the methodology spread across all natural sciences, namely epidemiology, ecology and biology. ABC is based on moments matching: the moments of the model are matched with the ones observed from the data. Moments are simulated according to the observed sample size and inference is based on the Euclidean distance between the simulated moments and the observed ones.

The use of moments of ABC-techniques makes the methodology similar to the GMM-style estimators: the Generalized Method of Moments (GMM, Hansen 1982) and its simulated version (the Simulated Method of Moments, SMM). In Section 3, a comparison between the ABC and a bayesian version of a GMM estimator

¹Gust et al. estimate a similar model using only three observables

(Kim,2002) is assessed. As it will become clear ABC estimators present a series of advantages with respect to the GMM-style estimators. ABC are particularly fit in the estimation of models whose likelihood computation is troublesome or whose moments distribution prevents the use of GMM-style estimators (irregular moments distribution, non-gaussian moments, short samples).

ABC methods have a Bayesian structure: the moments matching procedure updates a prior distribution to deliver an approximate posterior distribution. Approximation is a result of using the moments rather than computing the likelihood function of the model. The pseudo-algorithm by Pritchard (2000) clarifies the mechanism at the core of ABC methods and goes under the name of ABC-rejection:

- Draw θ_i from the prior distribution $p(\theta)$
- Simulate the model and get the variable $\mathbf{y}_i$
- Compute the summary statistics $\mathbf{s}_i$
- If the Euclidean distance $\rho||\mathbf{s}_i - \mathbf{s}|| < \epsilon$ accept θ_i otherwise reject it
- Repeat the procedure for N times

where $\mathbf{s}_i$ is the vector of moments from the simulated sample, $\mathbf{s}$ is the vector of moments of the observed data, ϵ is the tolerance level.

In other words, in ABC-rejection the model is simulated a number of times conditional on parameters drawn from the prior distribution. Moments from these simulations are computed and matched against the observed moments. For each simulation the Euclidean distance is computed. If the euclidean distance is smaller than a fixed threshold, the simulation is accepted. The parameters of the accepted simulations are a sample from the approximate posterior distribution.

The Bayes Rule of the Bayesian statistics is approximated:

$$P(\theta|\mathbf{y}) \propto L(\mathbf{y}|\theta)P(\theta) \rightarrow P(||\mathbf{s}_i - \mathbf{s}|| < \epsilon)P(\theta) \quad (1)$$

The likelihood function is approximated by the accept-reject step on the euclidean distances criterion. If the moments used in the estimation are sufficient statistics of the model, for $\epsilon \rightarrow 0$ and $N \rightarrow \infty$ the sequence of θ 's accepted converges to the posterior distribution. A large number of simulations needs to be run to reduce the error introduced by the simulation step. When the number of parameters to infer increases, so does the number of moments to use. The probability that the Euclidean distance is below the threshold is smaller and a larger number of simulations are run to obtain N accepted draws.

This brings to high inefficiency (the acceptance ratio gets small) and this may make simple ABC impractical (curse of dimensionality). A series of more sophisticated method has been developed to tackle the curse of dimensionality. Accepted simulations can be assigned a weight according to a kernel weighting function. The argument of the kernel is the euclidean distance: the smaller the distance, the larger the weight. In this paper, this method is called *ABC-kernel* and coincides with the simulated Bayesian version of the estimator proposed by Creel and Kristensen, 2012. In order of time, the ABC-rejection is the first method developed and is at the core of the other more sophisticated methods. ABC methods are mainly divided in three big subsets:

- ABC-regression;
- ABC-MCMC;
- ABC-SMC.

The three groups adopt different strategies to tackle the curse of dimensionality and the low efficiency of Pritchard algorithm. In particular the first solution runs a post-sampling correction on the accepted parameters, the last two draw parameters more efficiently.

2.1 ABC with local linear regression

ABC-rejection is affected by the curse of dimensionality: to estimate a large set of parameters, we need to increase the number of summary statistics in the Euclidean distance computation. The probability of the simulated parameters to be accepted decreases and a higher number of simulations have to performed. This may have a huge impact on the feasibility of the estimation procedure. Besides, to increase the tolerance level can strongly compromise the approximation of the posterior distribution due to a larger simulation error. ABC-regression increases the efficiency of ABC through a post-sampling correction.

Three main refinements are introduced after the accept-reject step:

- The moments are rescaled by their median absolute deviation: this transforms the previous rectangular acceptance region in a sphere.
- Each accepted simulation is assigned a weight according to its euclidean distance: the smaller the distance ρ_i , the larger the weight W_i . An Epanechnikov weighting function is generally used, but the algorithm is compatible with other kinds of kernel (normal, triangular and so forth). ².

²This correction coincides with the Indirect Likelihood Inference by Creel and Kristensen (2013)

- The accepted parameters are corrected exploiting the result of a regression run after the accept-reject (hence the name ABC-regression). Each parameter is updated according to the result of a local linear regression of the accepted parameters on the discrepancies between simulated moments and observed ones (Beaumont et al. (2002)).

In ABC regression (Beaumont, 2002), ABC is equivalent to a problem of conditional density estimation, where a joint density distribution $P(\mathbf{s}_i, \theta_i)$ is updated through an accept-reject algorithm:

$$P(\theta|\mathbf{s}) = \frac{p(\mathbf{s}_i, \theta)}{I \{ \rho|\mathbf{s}_i - \mathbf{s}| < \epsilon \}} \quad (2)$$

For this reason, conditional density estimation techniques (Fan and Gijbels, 1992) estimation are borrowed and incorporated in the ABC algorithms.

The ABC-regression pseudo-algorithm is:

- Draw θ_i from the prior $P(\theta)$;
- Simulate the model and obtain the observable variables $\mathbf{y}_i$;
- Compute the simulated moments $\mathbf{s}_i$ and the absolute standard deviation; for each moment k_j ;
- Compute the Euclidean distance for each simulation;

$$\rho|\mathbf{s}_i, \mathbf{s}| = \sqrt{\sum_{j=1}^s (\mathbf{s}_i/k_j - \mathbf{s}/k_j)^2} \quad (3)$$

- Select the tolerance level such that a fraction of the simulated parameters is accepted $P_\epsilon = N/M$;
- Each accepted draw is assigned a weight according to the Epanechnikov kernel:

$$K_\epsilon(\rho_i) = \begin{cases} \epsilon^{-1}(1 - (\frac{\rho_i}{\epsilon})^2) & \rho_i \leq \epsilon \\ 0, & \rho_i > \epsilon \end{cases}$$

;

- Apply a local linear regression to the linear model:

$$\theta_i = \alpha + (\mathbf{s}_i - \mathbf{s})'\beta + \epsilon_i, \quad (4)$$

for $i = 1, \dots, N$.

- Adjust the parameter given the results of the local linear regression:

$$\theta^* = \theta - (\mathbf{s}_i - \mathbf{s})' \hat{\beta}, \quad (5)$$

which is equivalent to compute: $\theta_i^* = \hat{\alpha} + \hat{\epsilon}_i$

The adjusted parameters associated to their kernel weights are random draws of the approximate posterior distribution.

The initial part of the ABC-regression is the simple ABC rejection. The accepted parameters are corrected given two assumptions on the relation between the parameters drawn and the summary statistics simulated:

- Local linearity: a local linear relationship between the discrepancies of the moments and the parameters holds in the vicinity of the observed moment s such that the parameters can be expressed by the following equation:

$$\theta_i = \alpha + (\mathbf{s}_i - \mathbf{s})' \beta + \epsilon_i; \quad (6)$$

- Errors ϵ_i 's have zero mean, are uncorrelated and homoskedastic.

In general, linearity only in the vicinity of $\mathbf{s}$ is a more palatable assumption than *global* linearity. In the local linear regression to estimate the coefficients for α and β , the minimized object is:

$$\sum_{i=1}^m \{\theta_i - \alpha - (\mathbf{s}_i - \mathbf{s})^T \beta\}^2 K_\delta(\|\mathbf{s}_i - \mathbf{s}\|) \quad (7)$$

In ABC literature, Epanechnikov kernel function is the one more common but others are feasible. In Eq.(7), the only difference with respect to the standard OLS is that the squared errors are weighted according to the distance ρ_i associated to the parameter θ_i . The solution is given by:

$$(\hat{\alpha}, \hat{\beta}) = (XWX)'(XW\theta) \quad (8)$$

Where $X = (\mathbf{s}_i - \mathbf{s})$ for $i = 1, \dots, N$ and W is a diagonal matrix, where each non zero element is $K_\delta(\|\mathbf{s}_i - \mathbf{s}\|)$.

The estimates for α and β are used in the adjustment step, through the adjustment equation 5.

In conditional density estimation terms: $E[\theta | \mathbf{s}_i = \mathbf{s}] = \alpha$.

The posterior mean coincides with the Nadaraya-Watson estimator (Nadaraya, 1964, Watson, 1964), as suggested by Blum and Francois (2010) :

$$\alpha = \frac{\sum_i \theta_i^* K_\delta(||\mathbf{s}_i - \mathbf{s}||)}{\sum_i K_\delta(||\mathbf{s}_i - \mathbf{s}||)} \quad (9)$$

Blum and Francois (2010) add further step: a *correction for heteroskedasticity* in the adjustment step with non-linear regression in lieu of the local linear regression. For the sake of simplicity, here the local linearity assumption is maintained allowing the variance of the errors to change with the moments (Beaumont, 2010). The heteroskedastic is:

$$\theta_i = \alpha + (\mathbf{s}_i - \mathbf{s})' \beta + \epsilon_i = \alpha + (\mathbf{s}_i - \mathbf{s})' \beta + \sigma_i * \xi_i, \quad (10)$$

where σ_i^2 is the variance of the error conditional on observing the simulated moments $Var[\theta|\mathbf{s}_i]$ and $\xi_i \sim N(0, 1)$.

In this new procedure (ABC-regression with correction for heteroskedasticity) estimates α and β remain the same while in a further step the conditional variance for each draw is estimated. Finally, the correction mechanism is applied.

Blum and Francois model the conditional variance on the moments discrepancy by a second local linear model, borrowing from Fan and Yao (1998). A second local linear regression is run and the conditional variance for each draw σ_i is estimated :

$$\log(\epsilon_i)^2 = \tau + (\mathbf{s}_i - \mathbf{s})' \pi + v_i, \quad (11)$$

where v_i is *iid* with mean zero and common variance.

In this second local linear regression, the following object is minimized:

$$\min \left\{ \log(\hat{\epsilon}_i)^2 - (\mathbf{s}_i - \mathbf{s})' \pi \right\} K_\delta(||\mathbf{s}_i - \mathbf{s}||) \quad (12)$$

where $\hat{\epsilon}_i$'s are the heteroskedastic errors estimated in the first regression.

The variance conditional on the observed moments is $\sigma^2 = Var[\theta|s]$ is obtained according to

$$\hat{\sigma} = \hat{\tau} \quad (13)$$

while the the variance conditional on each *simulated* moments is

$$\hat{\sigma}_i = \hat{\tau} + (\mathbf{s}_i - \mathbf{s})' \hat{\pi} \quad (14)$$

Values obtained in 14 are used in the new post-sampling correction equation 15 where the magnitude of each *heteroskedastic* error ϵ_i is corrected by the estimated standard deviation $\hat{\sigma}_i$:

$$\theta^* = \hat{\alpha} + \frac{\hat{\sigma}}{\hat{\sigma}_i} \hat{\epsilon}_i \quad (15)$$

When the associated variance is higher than the variance conditional on the observed moments, the ratio $\frac{\hat{\sigma}}{\hat{\sigma}_i}$ is lower than 1 and the magnitude of the correction will be decreased with respect to the estimated $\hat{\epsilon}_i$.

ABC-regression allows to increase the tolerance level (i.e. increase the fraction of accepted simulations), making the algorithm computationally more efficient. Nonetheless, when the dimensionality of the parameters increases, the algorithm can deliver unstable results. Besides, some problems in the adjustment step can arise when the local linearity assumption does not hold: when the observed moments lie at the boundary of the simulated moments, adjusted values can be updated outside the support of the prior distribution (extrapolating rather than interpolating). Some refinements have been found by the literature to fix this problem, but a general consensus has not been reached.

Before adopting ABC-regression, drawing scatter plots can be useful to assess the informativeness of the moments regard the parameters to infer. In particular, (local) linear relations between moments and parameters can be found. When the dimensionality of the problem makes both ABC-rejection and ABC-regression impractical, the ABC-SMC is the technique more fit to tackle the curse of dimensionality, as it will be shown in the final Section.

2.2 ABC-MCMC

ABC-MCMC methods draw parameters from a distribution closer to the posterior. This increases the acceptance rate of the algorithm. The algorithm developed by Marjoram et al. (2003) is the following:

- For $t = 0$, Draw $\theta \sim \pi(\theta)$;
- For $t \geq 1$ draw from:

$$\theta' \sim K(\theta|\theta^{t-1}); \quad (16)$$

- Simulate and produce the moments conditional on θ^t ;
- If $\rho(S(x), S(y)) < \epsilon$
 - Draw $u \sim U(0, 1)$,

– If

$$u \leq \frac{\pi(\theta')}{\pi(\theta)^{t-1}} \frac{K(\theta^{t-1}|\theta')}{K(\theta'|\theta^{t-1})} \quad (17)$$

then, $\theta^t = \theta'$; otherwise $\theta^t = \theta^{t-1}$

otherwise $\theta^t = \theta^{t-1}$

The MCMC produced by the algorithm is an approximation of the posterior distribution. Problems associated with ABC-MCMC are mainly related to presence of multimodality and mixing problems.

2.3 ABC-Sequential Montecarlo

ABC-SMC nests ABC into the structure of a SMC technique: the initial particles are drawn from a proposal distribution. Each particle is a vector of parameters. The distribution is iteratively updated to converge to the target distribution. At each step particles are perturbed according to a Kernel function. Each particle is accepted or rejected according to the Euclidean distance, choosing a decreasing tolerance level such that $\epsilon_t \leq \epsilon_{t-1}$. If accepted, the particle is assigned a weight taking into account the Kernel function. A resampling procedure is envisaged to avoid sample degeneracy (i.e. few particles ending up hoarding much of the weight). The algorithm is the following:

1. Initialize the tolerance level sequence: $\epsilon_1, \epsilon_2, \epsilon_3 \dots \epsilon_T$ and select a sampling distribution μ_i . Set the iteration indicator $t = 1$.
2. Set the particle indicator $i = 1$ and:
 - If $t = 1$, draw the swarm of particles $\{\theta_1 \theta_2 \dots \theta_N\}$ from the importance distribution μ_1 .
 - If $t > 1$, sample the new swarm $\{\theta_{i,t-1}^{**}\}_{i=1}^N$ with weights $\{W_{i,t-1}^{**}\}_{i=1}^N$ and perturb each particle according to a transition kernel $\theta^{**} \sim K_t(\theta|\theta^*)$
3. Simulate the model to obtain x^{**} conditional on each particle : if $\rho(S(x^{**}), S(x_0)) < \epsilon_t$ accept the particle, otherwise reject.
4. If accepted, assign the particle a weight:
 - If $t = 1$, $W_{i,1} = \frac{\pi(\theta_{i,1})}{\mu_1(\theta_{i,1})}$.
 - If $t > 1$,

$$W_{i,t} = \frac{\pi(\theta_{i,t})}{\sum_{j=1}^N W_{t-1}(\theta_{t-1,j}) K_t(\theta_{t,i}|\theta_{t-1,j})} \quad (18)$$

where $\pi(\theta)$ is the prior distribution for θ .

5. Normalize the weights such that $\sum_{i=1}^N W_{t,i} = 1$.
6. Compute the Effective Sample Size (ESS):

$$ESS = \left[\sum_{i=1}^N (W_{t,i})^2 \right]^{-1} \quad (19)$$

If the ESS is below $N\frac{1}{2}$, resample with replacement the particles according to the weights $\{W_{i,t}\}_{i=1}^N$ and obtain the new population with new weights $W_{t,i} = \frac{1}{N}$.

7. If $t < T$, return to (2).

This method does not get stuck in low probability areas or is able to explore the whole support also in case of multimodality. It eases the inefficiency in case of significant mismatch between prior and posterior. All these reasons make it particularly fit for the estimation of non-linear DSGE models.

3 A comparison with the Bayesian Limited Information Method

In this section the performance of the ABC estimators is compared with an increasingly popular alternative: the Limited Information Method (Kim, 2002). Its Bayesian version (the Bayesian Limited Information Method, henceforth BLI) is often interpreted as the Bayesian counterpart of the GMM-style estimators. The BLI is obtained by applying a Bayes Rule where the Prior distribution contains the extra data information and the likelihood is the joint probability of the *moments*, rather than of the data. Since the Central Limit Theorem applies, the likelihood is obtained relying on the asymptotic normal distribution of the moments (i.e. on the Central Limit Theorem).

Given the vector of parameters θ , the sample moments $\hat{\gamma}$ and the estimated variance of the moments $\hat{V}$, The Approximate Posterior distribution $P(\theta|\hat{\gamma}, \hat{V})$ is obtained according to the Bayesian updating rule:

$$P(\theta|\hat{\gamma}, \hat{V}) = \frac{P(\hat{\gamma}|\hat{\theta}, V)P(\theta)}{P(\hat{\gamma}|V)} \quad (20)$$

where T is the number of moments, $\hat{\gamma}$ is the vector of sample moments, $\gamma(\theta)$ is the vector of analytical moments depending on the parameter θ , $P(\theta)$ is the prior distribution. The likelihood $P(\gamma|\hat{\theta}, \hat{V})$, conditional on $\hat{V}$, is computed according to:

$$P(\gamma|\hat{\theta}, \hat{V}) = \frac{1}{(2\pi)^{(\frac{N}{2})}} |\hat{V}|^{-\frac{1}{2}} \exp \left\{ -\frac{T}{2} (\hat{\gamma} - \gamma(\theta))' V^{-1} (\hat{\gamma} - \gamma(\theta)) \right\}. \quad (21)$$

The role of moments and the Bayesian structure make the BLI the direct competitor of ABC estimators to check the small samples properties of the ABC estimator in a Bayesian framework. Interestingly, BLI can be interpreted as the Bayesian counterpart of the GMM and SMM estimators.

The relative performance of the ABC estimator with respect to the BLI method is measured in two Montecarlo exercises. The goal is to understand how much the presence of small samples and large persistence across the time series affect two estimators. The criteria for the comparison are twofold:

- the Root Mean Square Error (RMSE) with respect to the Full likelihood Posterior Mean;
- The Overlapping Ratio between the 90% Credible Intervals of the Approximate Posterior distributions and the Full Likelihood Posterior Distribution (our target distribution).

These two criteria analyse the estimators from a Bayesian perspective. RMSE measures how close are the two estimators to the Full likelihood Bayesian estimator (the Posterior Mean). The Overlapping Ratio captures which of the two methods deliver a better approximation of the posterior distributions. The RMSE is obtained by:

$$RMSE = \frac{1}{N} \frac{\sum (\hat{\theta}_{app} - \hat{\theta}_{full})^2}{\theta}, \quad (22)$$

where $\hat{\theta}_{app}$ is the mean of the posterior of one of the two approximating methods, $\hat{\theta}_{full}$ is the full likelihood posterior mean.

The Overlapping Ratio is obtained by:

$$OR = \frac{CI_{90\%, App} \cap CI_{90\%, Fl}}{CI_{90\%, App} \cup CI_{90\%, Fl}} \quad (23)$$

where $CI_{i-\%, App}$ is the i -th Percentile of the Approximate Posterior distribution, $\cap$ stands for Intersection and $\cup$ for Union. The Overlapping Ratio is always included in the interval $[-1, 1]$. For example if the two intervals perfectly coincide

the Overlapping Ratio equals 1, whereas if two degenerate posterior distributions do not overlap at all, the Overlapping Ratio equals -1.

The BLI estimator relies on the usual regularity assumptions of the GMM-style estimators. The normality assumption allows the GMM-style estimators to compute the likelihood of the moments focusing just on the first and the second moments of the moments distributions, and compute the quadratic objective function to update the prior. With ABC methodology, the moments distribution is studied by simulating the model, according to the observed sample size. The departure from the normality assumption and the kernel exploration of distribution delivers more reliable estimators than the GMM style estimators when dealing with small samples and highly persistent cases.

This result has been partially pointed out by Creel and Kristensen (2012) in their Indirect Likelihood Inference with which ABC shares the same intuition and similar asymptotic results. In a first step, the experiment is run on simple AR(1) model. In the remainder of this section focus is on a RBC model subject to some weak identification issues.

3.1 Case 1: AR(1)

Despite its simple structure, the AR(1) process reproduce different estimation issues. Moreover, most exogenous processes generating stochasticity in DSGE models are AR(1) processes exhibiting different kind of persistence (from low persistence processes to Unit Roots). The AR(1) model is estimated by varying the sample size and the persistence of the process, in order to check if and when an estimator exploring the simulated distribution of the moments (ABC) outperforms one relying on the normality assumptions and focusing on the GMM-style quadratic objective function (BLI).

The estimation for each AR(1) process is run 1000 times. The sample sizes are 100, 300, 1000 observations. The autocorrelation factor tuning the persistence can assume the following values $\phi = [0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 0.95, 0.99]$. Increasing the persistence of the process and decreasing the sample size should favour ABC estimators both in terms of RMSE and Overlapping Ratios. Vice versa, the gap between the RMSEs and the ORs of the two estimators is expected to close increasing the sample size and lowering the persistence. The moment in the matching procedure is the first order autocovariance. The Prior distribution is a Uniform prior $\sim U[0, 1]$. For the ABC, AR(1) is simulated 10000 times, the Euclidean distances between the observed autocovariance and the simulated ones are computed and sorted out to

select the first percentile of the distribution. The curse of dimensionality does not affect the estimation: 10000 simulations are enough to get rid of the error induced by the simulations since the moment is a scalar. For this reason, the correction step of the ABC-regression and the Kernel Weighting do not improve the estimation results upon the ABC-rejection procedure. Only results for ABC-regression are reported for sake of brevity.

For the Bayesian Limited Information method, the likelihood of the autocovariance is computed and the prior updated. The posterior distribution is studied with the Importance Sampling algorithm: as importance distribution the prior distribution is used and 10000 samples are drawn for each estimation. The estimated variance $\hat{V}$ to condition the likelihood (and the posterior distribution) is computed with two alternatives: the HAC Variance Covariance Estimator or a bootstrapping procedure. In the first case, the Newey-West estimator is computed, using a Bartlett Kernel and the bandwidth equal to $B(T) = \text{floor}(4 * (T/100)^{(2/9)})$, where T is the sample size.

An alternative method is inspired to the solution proposed in Christiano, Trebandt and Walentin, where the covariance matrix is estimated through a bootstrap step. In the latter case, a first step estimator is computed to minimize a quadratic objective function using the identity matrix as variance covariance matrix. Afterwards, the AR(1) process is simulated for 1000 times (bootstrapping) to compute the autocorrelation for each bootstrap and the covariance of the moments $\hat{V}$ to compute the likelihood. Since working with one moment, the identity matrix at the initial step is simply the unity scalar and the covariance matrix is the covariance of the autocovariances computed in the bootstrap step.

Before exposing the results of the experiment, it can be interesting to give a quick look to the distribution of the autocovariances obtained by simulating the AR(1) process using different autocorrelations (from low persistency up to almost unit roots). In Fig. 1, the distribution of the sample autocovariances when $\phi = 0.5$ is reported. The sample size varies from 50 to 1000 observations. The distribution of the autocovariances converge quickly to a Normal distribution with the mean of the population autocovariance ($\gamma = \phi/(1 - \phi^2)$), represented by the pink plane. In the the highly persistent case (Fig.2) when $\phi = 0.99$, the convergence to the normal distribution is much slower and even with a sample size of 5000 observations, the distribution of the autocovariance is skewed and not centred around the population autocovariance. These results render a simple intuition on the expected (and found) results in the Montecarlo experiments. Fig. 3, Fig. 4, and Fig. 5 show the evolution of the RMSEs with respect to the Full likelihood posterior mean varying

the persistence from $\phi = 0.1$ up to $\phi = 0.99$. The three figures are generated using different sample size: respectively 100, 300 and 1000 observations. Comparing the three figures, the gap between the two methods is larger in favour of ABC in small samples and reduces increasing the sample size. Moreover, for each sample size, increasing the persistence widens the gap in favour of ABC. Among the different approaches to estimate the variance covariance matrix of the moments $\hat{V}$, the HAC Newey-West estimator ensures smaller RMSEs especially in highly persistent cases and small samples, while the bootstrapping methods has smaller RMSE with low autocorrelations. In large samples, the RMSEs converge, at least up to $\phi = 0.95$. These results were widely expected in the light of the distributions juxtaposed in Figs. 1,2. Figs. 6 7 and 8 show instead the evolution of the Overlapping Ratios passing from low autocorrelations to almost unit roots.

Again, the samples are made of 100, 300 and 1000 observations. Our intuition is confirmed by the results: the OR gap between ABC methods and BLI is larger in general for highly persistent processes proving that ABC outperforms BLI in approximating the posterior distributions under certain conditions: the smaller the sample size, the larger the gap between the methods in favour of ABC. Also from this standpoint, results suggest that among the BLI estimators, the HAC estimation of the variance covariance matrix has a larger OR values than the Bootstrapping Procedure for persistent processes, while the opposite is true for the low persistent cases.

3.2 Case 2: A RBC with identification issues

In this second section, the performance of the two estimators is studied in a more complex and real world application. The experiment is run on a linear RBC model with three structural shocks and three observables. Again the comparison is run from a Bayesian perspective, taking the Full likelihood posterior distribution as reference and trying to capture to which extent the two approximate posterior distributions approximate the results of the true full likelihood posterior distributions.

The RBC estimated by Creel and Kristensen (2012) is a plain-vanilla RBC model with no problems of identification, given the simple structure of the model with just one structural shock on productivity. The RBC studied in this section encounters some identification issues concerning the preference parameters, due to the presence of three stochastic processes: a productivity shock on the production function, a shock on the preference affecting the labour supply and a shock on the interest rate requested by the household. The presence of these three shocks permit to estimate the full likelihood distribution using three observable variables without the need for

measurement errors.

The households maximize the following expected sum of the utility functions:

$$\max E_t \left(\sum_{t=0}^{\infty} \beta^t \left(\ln C_t - B_t \frac{H_t^{1+\frac{1}{\nu}}}{1+\frac{1}{\nu}} \right) \right) \quad (24)$$

subject to the budget constraint:

$$C_t + I_t = W_t H_t + D_t R_t K_t. \quad (25)$$

E_t stands for the expectation operator, C_t is the consumption, H_t are the hours offered by each household, B_t is the shock to the preference (namely the labour supply) (Rios-Rull et al., 2012) and D_t is the shock to the interest rate requested by the household like in Smets and Wouters (2007). β is the subjective discount factor and ν is the Frisch elasticity. Capital K_t is cumulated according to the following rule:

$$K_{t+1} = (1 - \delta)K_t + I_t \quad (26)$$

where δ is the depreciation rate and I_t is the investment. Firms choose how much capital and hours to employ in the production function given the technology A_t :

$$Y_t = A_t K_t^\alpha H_t^{1-\alpha} \quad (27)$$

The market clearing is defined by:

$$Y_t = C_t + I_t \quad (28)$$

The economy is subject to the following three structural shocks:

$$\log(A_{t+1}) = \rho_a \log(A_t) + \sigma_a \epsilon_a \quad (29)$$

$$\log(B_{t+1}) = \rho_b \log(B_t) + \sigma_b \epsilon_b \quad (30)$$

$$\log(D_{t+1}) = \rho_d \log(D_t) + \sigma_d \epsilon_d \quad (31)$$

The technology shock A_t is the standard shock of the RBC literature (Kydland and Prescott, 1983). The shock on the preferences B_t perturbs the labour supply hitting the marginal rate of substitution between consumption and leisure (see Rios-Rull, 2012). The shock on D_t is a shock on the interest rate requested by the households and can be interpreted as a shock to the risk premium. (Smets and Wouters, 2007).

The model equilibrium is obtained by the following equations:

$$H_t = \left(\frac{1}{B_t} \frac{W_t}{C_t} \right)^\gamma \quad (32)$$

$$\frac{1}{C_t} = \beta \left(\frac{1}{C_{t+1}} ((1 - \delta) + D_{t+1} R_{t+1}) \right) \quad (33)$$

$$Y_t = C_t + I_t \quad (34)$$

$$K_{t+1} = K_t(1 + \delta) + I_t \quad (35)$$

$$R_t = \alpha A_t K_t^{\alpha-1} H_t^{1-\alpha} \quad (36)$$

$$W = (1 - \alpha) A_t K_t^\alpha H_t^{-\alpha} \quad (37)$$

$$Y_t = A_t K_t^\alpha H_t^{1-\alpha} \quad (38)$$

Eq. 32 is the intratemporal choice between consumption and leisure, Eq. 33 is the Euler Equation. Eqs. 34 and 35 are the resource constraints and market clearing conditions completing the equilibrium of the model. Eq. 35 is the law of motion of capital and Eqs. 36, 37, 38 are the exogenous processes.

The experiment adopts an informative prior distribution of the same fashion that Rios-Rull et al.(2012) use to estimate a state-of-the-art Real Business Cycle. Informativeness in the prior distribution eases the identification issues associated to the preferences parameters.

Each Montecarlo experiment is made of 100 repetitions. The RMSE and the Overlapping Ratio are computed using different sample sizes: 100, 200, 500 observations. The data generating parameters are the following: $\beta = 0.95$, $\gamma = 2$, $\rho_a = 0.95$, $\rho_b = 0.95$, $\rho_d = 0.95$, $\sigma_a = 0.01$, $\sigma_b = 0.01$, $\sigma_d = 0.01$. The persistence of the processes plays in favour of the ABC-estimators, especially in light of the results previously obtained in the AR(1) case. The moments are the covariances and the first order autocovariances of three observables: income Y_t , hours H_t and investments I_t .

The prior distribution is indicated in Table 1. Concerning the ABC methods, RBC is simulated 5000 times, the tolerance level is such that the acceptance ratio of the simulations is equal to 5%. The results of ABC-rejection, ABC-kernel, ABC-regression and ABC-regression+HC, ABC-OLS (ABC-regression where the regression is simply linear) are reported.

The variance covariance matrix of the BLI estimator is obtained through the HAC Newey-West estimator. For each Full likelihood and BLI estimation, a MCMC is drawn following the steps listed in An and Shorfheide (2007). Each chain contains

10000 draws with a burn-in period of 1000 draws. Table 2 contains the results of the RMSE for the case of 100 observations, informative prior and high persistence of the process. ABC RMSEs are smaller than the BLI RMSEs. Tables 3 and 4 report the RMSEs respectively for 200 and 500 observations. The gap between the estimators is still in favour of the ABC.

Overlapping Ratios of the 90% credible intervals of the approximate posterior distributions and the Full likelihood posterior distribution are compared. The results are respectively reported in Tables 5,6,7. ABC outperforms BLI method in approximating the full likelihood posterior distribution under the three different sample sizes: 100, 200 and 500 observations.

4 An application to the Zero Lower Bound

The financial crisis of 2008, the Great Recession and the years of slow growth represent a big challenge to DSGE modelling and their estimation. Since the beginning of the crisis and with different timings, many central banks lowered interest rates at their minimum and maintained them there for more than 5 years. In 2014Q4, interest rates in U.S, Euro Area, U.K., Sweden and other economic areas are still at the zero lower bound. From a model perspective, the binding constraint on the policy rate and the gap between the Taylor rule implied interest rate and the actual one cause a non-linearity to take into account in the model solution.

In this section, ABC-SMC is applied on the estimation of a newkeynesian model with an occasional binding constraint on the zero lower bound. The non-linear model is borrowed by Fernandez-Villaverde et al. (2012). The notation is the same for the sake of comparison between the calibrated values of the original papers and the estimates of this section. As it appears clear from Section 3, ABC techniques study the simulated distribution of the moments without relying on the normality and regularity assumptions made in the GMM-style estimators. Moreover, in ABC non gaussian moments can be exploited (i.e. binomial, multinomial etc.) Here, the following moments are used in the estimation together with the usual covariances:

- Frequency of the zero lower bound, number of episodes at the zero lower bound in the sample;
- Frequency of recession events, number of recession episodes;
- Frequency of deflation events, number of deflation episodes.

In this paper the model is solved according to the Piecewise linear solution, using the MATLAB routine provided by Guerrieri and Iacoviello (2014). The piecewise linear solution is quick to obtain with respect to the other non-linear methods and can handle medium-size DSGE models. This allows to obtain a large numbers of simulations in short range of time. The sample includes six observable variables and includes observations up to 2014Q3. The main exercise uses data starting for the beginning of the Great Moderation (1983Q1). Hence in the sample, the ZLB binds for more than one fifth of the sample. This, together with the use of non-conventional features of the data (frequency of the ZLB and so forth), tries to capture the effects of the exit from the Great Moderation in the estimate results.

4.1 The model

The model is a standard newkeynesian model with occasionally positivity constraint on the interest rate. A household maximizes her utility consuming and providing labour (the unique productive factor) to intermediate firms operating in monopolistic competition, readjusting prices according to Calvo type of contracts. The differentiated products are then assembled by retail firms operating in perfect competition.

Households maximise the following utility function separable in consumption c_t and labour l_t .

$$\sum_{i=0}^{\infty} \left(\prod_{i=0}^t \beta_i \right) \left\{ \log c_t - \psi \frac{l_t^{1+\phi}}{1+\phi} \right\} \quad (39)$$

where ϕ is the inverse of the Frisch labour supply elasticity and β_t is the subjective discount factor subject to stochastic fluctuations around the mean β :

$$\beta_{t+1} = \beta^{1-\rho_b} \beta_t^{\rho_b} \exp(\sigma_b \epsilon_{b,t+1}) \quad (40)$$

with $\epsilon_{b,t+1} \sim N(0, 1)$. ρ_b and σ_b are respectively the autocorrelation and the standard deviation of the AR(1) process. The household maximizes her utility subject to the budget constraint:

$$c_t + \frac{b_{t+1}}{p_t} = w_t l_t + R_{t-1} b_t / p_t + T_t + F_t \quad (41)$$

where b_t is a nominal government bond that pays a nominal interest rate R_t . p_t is the price level, whereas T_t and F_t are respectively the lump sum taxes and the profits of the firms. Retail firms reassemble intermediate goods y_{it} and the technology:

$$y_t = \left(\int_0^1 y_{it}^{\frac{\epsilon-1}{\epsilon}} di \right)^{\frac{\epsilon}{\epsilon-1}} \quad (42)$$

with ϵ is the elasticity of substitution. Final producers maximize their profit taking into account intermediate goods prices p_{it} , final prices p_t . The demand for each good will follow:

$$y_{it} = \left(\frac{p_{it}}{p_t} \right)^{-\epsilon} y_t, \quad (43)$$

and the price of the final good will be equal to:

$$p_t = \left(\int_0^1 p_{it}^{1-\epsilon} di \right)^{\frac{1}{1-\epsilon}}. \quad (44)$$

The wholesale firms operate according to the production function:

$$y_{it} = A_t l_{it}, \quad (45)$$

where the productivity A_t evolves according to the law of motion:

$$A_t = A^{1-\rho_A} A_{t-1}^{\rho_A} \exp(\sigma_A \varepsilon_{A,t}) \quad (46)$$

with $\varepsilon_t \sim N(0, 1)$. The marginal costs are $mc_t = \frac{w_t}{A_t}$.

The firms choose their price according to a Calvo rule, where each period just a fraction $1 - \theta$ firms can re-optimize their prices p_{it} . Firms will choose their price to maximize the profits:

$$\max_{p_{it}} E_t \sum_{\tau=0}^{\infty} \theta^\tau \left(\prod_{i=0}^{\tau} \beta_{t+1} \right) \frac{\lambda_{t+1}}{\lambda_t} \left(\frac{p_{it}}{p_{t+\tau}} - mc_{t+\tau} \right) y_{it+\tau} \quad (47)$$

s.t.

$$y_{it} = \left(\frac{p_{it}}{p_t} \right)^{-\epsilon} y_t \quad (48)$$

where λ_{t+s} is the Lagrangian multiplier for the household in period $t + s$. Two auxiliary $x_{1,t}$ and $x_{2,t}$ are used to define the solution to the maximization problem:

$$\epsilon x_{1,t} = (1 - \epsilon x_{2,t}) \quad (49)$$

$$x_{1,t} = \frac{1}{c_t} mc_t y_t + \theta E_t \beta_{t+1} \Pi_{t+1}^\epsilon x_{1,t+1} \quad (50)$$

$$x_{2,t} = \frac{1}{c_t} \Pi_t^* y_t + \theta E_t \beta_{t+1} \Pi_{t+1}^{\epsilon-1} \frac{\Pi_t^*}{\Pi_{t+1}^*} x_{2,t+1} = \Pi_t^* \left(\frac{1}{c_t} y_t + \theta E_t \beta_{t+1} \frac{\Pi_{t+1}^{\epsilon-1}}{\Pi_{t+1}^*} x_{2,t+1} \right) \quad (51)$$

where $\Pi_t^* = \frac{p_t^*}{p_t}$. Inflation dispersion will be equal to:

$$1 = \theta \Pi_t^{\epsilon-1} + (1 - \theta) (\Pi_t^*)^{1-\epsilon}. \quad (52)$$

The government sets the nominal interest rate:

$$R_t = \max [R_t, 1], \quad (53)$$

with the notional interest rate Z_t :

$$Z_t = R^{1-\rho_r} R_{t-1}^{\rho_r} \left[\left(\frac{\Pi_t}{\Pi} \right)^{\phi_\pi} \left(\frac{y_t}{y_{t-1}} \right)^{\phi_y} \right]^{1-\rho_r} m_t \quad (54)$$

with m_t being the monetary policy *iid* shock $m_t = \exp(\varepsilon_{m,t} \sigma_m)$, $\varepsilon_{m,t} \sim N(0, 1)$. The gross interest rate is equal to the notional interest rate as long it is larger than 1, since it cannot be set below 1 (the zero lower bound, ZLB).

The government sets also the spending:

$$g_t = s_{g,t} y_t \quad (55)$$

$$s_{g,t} = s_g^{1-\rho_g} s_{g,t-1}^{\rho_g} \exp(\sigma_g \varepsilon_{g,t}) \quad (56)$$

with $\varepsilon \sim N(0, 1)$. Since the agents are ricardian, we can set $b_t = 0$.

After aggregation we obtain:

$$y_t = \frac{A_t}{v_t} l_t \quad (57)$$

with v_t is the loss of efficiency introduced by the price dispersion:

$$v_t = \int_0^1 \left(\frac{p_{i,t}}{p_t} \right)^{-\epsilon} di \quad (58)$$

Moreover, following the Calvo pricing properties we can write:

$$v_t = \theta \Pi_t^\epsilon v_{t-1} + (1 - \theta) (\Pi_t^*)^{-\epsilon}. \quad (59)$$

4.2 The Equilibrium

The Equilibrium is given by the sequence

$$\{y_t, c_t, l_t, mc_t, x_{1,t}, x_{2,t}, w_t, \Pi_t, \Pi_t^*, v_t, R_t, Z_t, \beta_t, A_t, m_t, g_t, b_t, s_{g,t}\}_{t=0}^\infty \quad (60)$$

. The equilibrium is defined by the following equations.

The intertemporal and the intratemporal household F.O.Cs:

$$\frac{1}{c_t} = E_t \left\{ \frac{\beta_{t+1}}{c_{t+1}} \frac{R_t}{\Pi_{t+1}} \right\}, \quad (61)$$

$$\psi l_t^\phi c_t = w_t \quad (62)$$

The solution of the maximization problem of the firms:

$$mc_t = \frac{w_t}{A_t}, \quad (63)$$

$$\epsilon x_{1,t} = (1 - \epsilon x_{2,t}), \quad (64)$$

$$x_{1,t} = \frac{1}{c_t} mc_t y_t + \theta E_t \beta_{t+1} \Pi_{t+1}^\epsilon x_{1,t+1}, \quad (65)$$

$$x_{2,t} = \frac{1}{c_t} \Pi_t^* y_t + \theta E_t \beta_{t+1} \Pi_{t+1}^{\epsilon-1} \frac{\Pi_t^*}{\Pi_{t+1}^*} x_{2,t+1} = \Pi_t^* \left(\frac{1}{c_t} y_t + \theta E_t \beta_{t+1} \frac{\Pi_{t+1}^{\epsilon-1}}{\Pi_{t+1}^*} x_{2,t+1} \right). \quad (66)$$

The government equations are:

$$R_t = \max [R_t, 1], \quad (67)$$

$$Z_t = R^{1-\rho_r} R_{t-1}^{\rho_r} \left[\left(\frac{\Pi_t}{\Pi} \right)^{\phi_\pi} \left(\frac{y_t}{y_{t-1}} \right)^{\phi_y} \right]^{1-\rho_r} m_t. \quad (68)$$

Inflation evolution and price dispersion:

$$1 = \theta \Pi_t^{\epsilon-1} + (1 - \theta) (\Pi_t^*)^{1-\epsilon}, \quad (69)$$

$$v_t = \theta \Pi_t^\epsilon v_{t-1} + (1 - \theta) (\Pi_t^*)^{-\epsilon}. \quad (70)$$

Market clearing conditions:

$$y_t = c_t + g_t, \quad (71)$$

$$y_t = \frac{A_t}{v_t} l_t. \quad (72)$$

The stochastic processes are:

$$\beta_{t+1} = \beta^{1-\rho_b} \beta_t^{\rho_b} \exp(\sigma_b \epsilon_{b,t+1}), \quad (73)$$

$$A_t = A^{1-\rho_A} A_{t-1}^{\rho_A} \exp(\sigma_A \epsilon_{A,t}), \quad (74)$$

$$s_{g,t} = s_g^{1-\rho_g} s_{g,t-1}^{\rho_g} \exp(\sigma_g \epsilon_{g,t}), \quad (75)$$

$$m_t = \exp(\epsilon_{m,t} \sigma_m). \quad (76)$$

4.3 Solution method

Standard perturbation methods provide local solutions and cannot handle models with occasionally binding constraint without adaptation. For this reason, Fernandez-

Villaverde et al. solve the model using a fully non-linear solution. In this paper, the model is solved by the piecewise linear solution method presented in Guerrieri and Iacoviello 2013. The routine codes are directly provided by the authors. Here the solution technique is shortly presented. For a more detailed exposition, see the original paper.

The piecewise solution method delivers a first order perturbation solution in a piecewise fashion. The presence of the occasionally binding constraint creates two regimes: in one the constraint is slack, in the other it is binding. In the current exercise, the unconstrained case is the reference regime, the constrained (ZLB binding) is the alternative. The solution that we obtain is not just the juxtaposition of two linear solutions: the policy coefficients depend on how long the regime is expected to last. How long the model lasts is influenced by the state vector. This feedback effect can produce an important non-linearity. The piecewise linear solution allows to obtain a large number of simulations and tackle the curse of dimensionality generated by dealing with solving non-linearly medium-scale economic models.³

To solve the model two conditions must hold:

- Blanchard-Khan conditions must hold in the reference regime;
- If the shocks hitting the economy take the model away from the reference regime to the alternative regime, in absence of future shocks the model must return to the reference regime.

For further details on the solution method see Guerrieri and Iacoviello (2014).

4.4 Estimation strategy

The model is estimated using six quarterly macroeconomic time series for the US economy, taken from FRED dataset and used as observable variables: the log difference of Real GDP per person, the log difference of Real Consumption, log hours worked, the log GDP deflator, the log difference of real wage, the log FED funds rate. The observable equations are the following:

$$\log \Delta GDP_t = 100(y_t - y_{t-1}) + \gamma, \quad (77)$$

$$\log \Delta CONS_t = 100(c_t - c_{t-1}) + \gamma, \quad (78)$$

$$\log \Delta WAGES_t = 100(w_t - w_{t-1}) + \gamma, \quad (79)$$

³A drawback of this solution method is that it assumes that agents do not expect future shocks hitting the economy in the following periods. Hence precautionary savings are not considered.

$$\log HOURS_t = 100l_t + \bar{l}, \quad (80)$$

$$\log \Delta DEF L_t = 100 * (p_t * \pi + \pi - 1), \quad (81)$$

$$\log FEDDUNDS_t = 100(\exp(r_t) * RSS - 1) - 1. \quad (82)$$

Where Δ is the difference operator, $RSS = \frac{\pi}{\beta}$. The type of dataset is very similar to the one used by Smets and Wouters, except for the investment that is excluded. As in Smets and Wouters, measurement errors are not necessary to estimate the model, differently from the particle filter case.

The ZLB period started in 2008Q3 up to the end of the sample. Different estimation exercises are performed. First, estimation is conducted according to four different time ranges:

- *Baseline*: the sample size spans from 1966Q1 to 2014Q3. It is the largest sample, it contains 185 observations and starts from the same quarter used in the main estimation exercise in Smets and Wouters (2003);
- *Great Moderation without the ZLB*: the dataset range goes from 1983Q1 until 2007Q4. The sample contains 96 observations and stops before the interest rate enters the Great Recession and hits the ZLB period;
- *Great Moderation and the Great Recession*: the sample spans from 1983 to 2014Q3 (125 observations). The economy is at the ZLB for approximately one fifth of the time. The final part of the sample contains the Great Recession and the slow recovery.
- *The Great Volatility II*: the sample spans from 2001Q1 until 2014Q3 (57 observations). The economy is at the ZLB for approximately 40% of the sample.

In a first case, the estimation is performed using just the covariances and the variances of the observable variables. The moments are computed and matched conditional on the two different regimes. In a second exercise, the estimation is performed using covariances and non-gaussian distributed moments:

- the frequency of being on the zero lower bound over the sample, the number of periods at the ZLB;
- the frequency of being in recession over the sample, the number of periods at of recession;
- the frequency of being in deflation over the sample, the number of periods at of deflation.

The priors used are common in literature and are listed in Table 8. ABC-SMC procedure is applied, until convergence of the posterior distribution. The model is simulated for 30000 times at the first iteration. In the first iteration, the 5% of the simulations is accepted according to the Euclidean distance. After the first iteration, each swarm of particle contains 1500 simulations and the particles are perturbed according to the kernel $K(\theta_{i,t}^{**}|\theta_{i,t}^*)$:

$$\theta_{i,t}^{**} \sim N(\theta_{i,t-1}^*, c * \Sigma), \quad (83)$$

where Σ is a diagonal matrix with the variances of the first iteration accepted parameters, scaled by a scalar $c = 0.02$, to tune the acceptance rate in the following iterations.

When $t = 1$, weights are assigned according to:

$$W_{i,1} = \frac{\pi(\theta_{i,1})}{\mu_1(\theta_{i,1})} = 1 \quad (84)$$

since the prior distribution $\pi(\theta)$ and the proposal $\mu_1(\theta_{i,1})$ coincide.

When $t \geq 2$ weights are assigned according to:

$$W_{i,t} = \frac{\pi(\theta_{i,t})}{\sum_{j=1}^N W_{t-1}(\theta_{t-1,j}) K_t(\theta_{t,i}|\theta_{t-1,j})}, \quad (85)$$

where $W_{i,t}$ is the weight of particle i at iteration t . $K_t(\theta_{t,i}|\theta_{t-1,j})$ is the kernel of the perturbation step. Particles of parameters are resampled when the effective sample size is smaller than 750 (half of the accepted sample obtained after first accept-reject).

The tolerance level is decreased of 0.1% at each iteration.

The convergence for the approximate posterior distribution can be intuitively checked confronting the different approximate posterior distribution obtained at each iteration. In this case, ten iterations are enough to insure convergence of the posterior distributions (Fig. 9).

4.5 Estimation results.

Results for the four different samples are reported in Tables 9-12. Estimate results are standard. Across the different sub-samples, standard deviations are larger for the GM+ZLB and the GV-II periods and smaller for the GM period. Moreover, autocorrelations for the preference AR(1) process and the TFP AR(1) process are

larger for GM+ZLB and the GV-II periods. Autocorrelation for the government spending process is larger for the baseline and the GM period. Concerning the monetary policy, no significant differences emerge, except for the autocorrelation of the interest rate, which is smaller in the GM period compared to the other subsamples estimates. To check how these estimates affect the dynamics of the model, for each subsample estimate the model is shocked for two consecutive periods by two preference shocks in a row. The magnitude of each shock is equal to two standard deviations of the shock. Such shocks send the model onto the Zero lower bound in all the examples, except for the Great Moderation period. As expected, the latter case is the one where variables move more moderately (Fig.10). The largest variations are found in the GV II case. These results suggest that even if the sub-samples differ only for small fractions of data, the estimation results provided evidence for different behaviour of the main variables.

If the non-gaussian moments are included estimates do not vary much compared to the case with only gaussian moments. This is probably attributed to the fact that the amount of new information introduced with the non gaussian moments is relatively small compared to the one provided by the moments already in use. Moreover, these moments seem to concern more the parameters affecting the steady state value of inflation and interest rates (the inflation target Π and the subjective discount factor β). As a result, the impulse responses look very similar (Fig.11). The GM period is the one with the smallest reactions to the shock. With respect to the case with only gaussian moments, except for the steady state values for interest rate and inflation which are lower. From this simple exercise, the non gaussian moments used (duration of ZLB, frequency of the ZLB, duration of deflation and so forth) appear to be useful to improve the efficiency in the identification of steady state parameters. A further investigation is required on this topic.

5 Conclusion

In this paper, Approximate Bayesian Computation techniques have been applied to the estimation of economic models. Two Montecarlo experiments have been assessed to analyse the small samples properties of ABC techniques. ABC performance is compared to the one of the Bayesian Limited Information Method (BLI), Kim (2002). BLI can be interpreted as a Bayesian version of GMM-style estimators.

ABC outperforms BLI both using an AR(1) at different persistence and an RBC model with large persistence. The performance is analysed through the lens of

Bayesian criteria:

- The RMSE with respect to the Full likelihood posterior mean;
- The Overlapping ratio between the approximate posteriors and the full likelihood posterior distribution.

This result holds stronger when dealing with small sample and large persistence data generating processes: ABC does not automatically rely on the normality assumption made on the moments. ABC explores the whole moments distribution. Other estimation exercises are provided. ABC-rejection and ABC-regression are applied to a vanilla RBC model. A newkeynesian model with occasionally binding constraint is applied. The model is solved and simulated using the piecewise linear approximation by Guerrieri and Iacoviello 2013. ABC-SMC is applied to tackle the curse of dimensionality. The model is estimated with different sub-samples. Results show different behaviours of the main variables according to the sub-sample estimates used to get the dynamics of the model. Great Moderation impulse responses strongly differ from the ones obtained using estimates which take into account the Great Recession and the ZLB period. Estimation is performed also using non-gaussian moments, as the frequency of hitting the zero lower bound or the number of ZLB episodes. In this case, inference is affected by using these unconventional features of the data, contributing to increase the efficiency in the identification of some parameters.

References

- [1] An, Sungbae, and Frank Schorfheide. "Bayesian analysis of DSGE models." *Econometric reviews* 26.2-4 (2007): 113-172.
- [2] Andreasen, Martin M., Jess Fernndez-Villaverde, and Juan Rubio-Ramrez. The Pruned State-Space System for Non-Linear DSGE Models: Theory and Empirical Applications. No. w18983. National Bureau of Economic Research, 2013.
- [3] Barro, Robert J. "Rare disasters and asset markets in the twentieth century." *The Quarterly Journal of Economics* 121.3 (2006): 823-866.
- [4] Beaumont, Mark A., Wenyang Zhang, and David J. Balding. "Approximate Bayesian computation in population genetics." *Genetics* 162.4 (2002): 2025-2035.
- [5] Beaumont, Mark A. "Approximate Bayesian computation in evolution and ecology." *Annual Review of Ecology, Evolution, and Systematics* 41 (2010): 379-406.
- [6] Benigno, Gianluca, Pierpaolo Benigno, and Salvatore Nistico. "Second-order approximation of dynamic models with time-varying risk." *Journal of Economic Dynamics and Control* (2013).
- [7] Blum, Michael GB, and Olivier Franois. "Non-linear regression models for Approximate Bayesian Computation." *Statistics and Computing* 20.1 (2010): 63-73.
- [8] Calvet, C., and Veronika Czellar. Accurate Methods for Approximate Bayesian Computation Filtering. *Technical Report*, HEC Paris, 2012.
- [9] Canova, Fabio, and Luca Sala. "Back to square one: identification issues in DSGE models." *Journal of Monetary Economics* 56.4 (2009): 431-449.
- [10] Creel, Michael, and Dennis Kristensen. "Indirect Likelihood Inference (revised)". No. 931.13. *Unitat de Fonaments de l'Analisi Economica (UAB) and Institut d'Analisi Economica (CSIC)*, 2013.
- [11] Christiano, Lawrence, Martin Eichenbaum, and Sergio Rebelo. When is the government spending multiplier large?. No. w15394. National Bureau of Economic Research, 2009.
- [12] Csillery, Katalin, et al. "Approximate Bayesian computation (ABC) in practice." *Trends in ecology and evolution* 25.7 (2010): 410-418.

- [13] Csillry, Katalin, Olivier Franois, and Michael GB Blum. "abc: an R package for approximate Bayesian computation (ABC)." *Methods in ecology and evolution* 3.3 (2012): 475-479.
- [14] Del Moral, Pierre, Arnaud Doucet, and Ajay Jasra. "An adaptive sequential Monte Carlo method for approximate Bayesian computation." *Statistics and Computing* 22.5 (2012): 1009-1020.
- [15] Del Negro, Marco, and Frank Schorfheide. "Forming priors for DSGE models (and how it affects the assessment of nominal rigidities)." *Journal of Monetary Economics* 55.7 (2008): 1191-1208.
- [16] Doucet, Arnaud, Nando De Freitas, and Neil Gordon. Sequential Monte Carlo methods in practice. Vol. 1. New York: Springer, 2001.
- [17] Eggertsson, Gauti B., and Michael Woodford. Optimal monetary policy in a liquidity trap. No. w9968. National bureau of economic research, 2003.
- [18] Fan, Jianqing, and Qiwei Yao. "Efficient estimation of conditional variance functions in stochastic regression." *Biometrika* 85.3 (1998): 645-660.
- [19] Fan, Jianqing, and Irene Gijbels. "Variable bandwidth and local linear regression smoothers." *The Annals of Statistics* (1992): 2008-2036.
- [20] Jess Fernndez-Villaverde, 2010. "The econometrics of DSGE models," *SERIEs, Spanish Economic Association*, vol. 1(1), pages 3-49, March.
- [21] Jesus Fernandez-Villaverde and Juan F. Rubio-Ramirez, 2006. "Estimating Macroeconomic Models: A Likelihood Approach," *NBER Technical Working Papers* 0321, National Bureau of Economic Research, Inc.
- [22] Fernndez-Villaverde, Jess, et al. Nonlinear adventures at the zero lower bound. No. w18058. National Bureau of Economic Research, 2012.
- [23] Geweke, John. "Using simulation methods for Bayesian econometric models: inference, development, and communication." *Econometric Reviews* 18.1 (1999): 1-73.
- [24] Geweke, John, and Hisashi Tanizaki. "Bayesian estimation of state-space models using the MetropolisHastings algorithm within Gibbs sampling." *Computational statistics and data analysis* 37.2 (2001): 151-170.

- [25] Geweke, John. "Bayesian comparison of econometric models." Federal Reserve bank of Minneapolis working paper 532 (1994).
- [26] Gordon, Neil J., David J. Salmond, and Adrian FM Smith. "Novel approach to nonlinear/non-Gaussian Bayesian state estimation." IEE Proceedings F (Radar and Signal Processing). Vol. 140. No. 2. IET Digital Library, 1993.
- [27] Ireland, Peter N., 2003. "Endogenous money or sticky prices?," *Journal of Monetary Economics*, Elsevier, vol. 50(8), pages 1623-1648, November.
- [28] Ireland, Peter N., 2004. "Money's Role in the Monetary Business Cycle," *Journal of Money, Credit and Banking*, Blackwell Publishing, vol. 36(6), pages 969-83, December.
- [29] Ireland, Peter N., 2004. "A method for taking models to the data," *Journal of Economic Dynamics and Control*, Elsevier, vol. 28(6), pages 1205-1226, March.
- [30] Andrews, Isaiah, and Anna Mikusheva. "Weak Identification in Maximum Likelihood: A Question of Information."
- [31] Andrews, Isaiah, and Anna Mikusheva. "Maximum likelihood inference in weakly identified DSGE models." (2011).
- [32] Judd, Kenneth L. Numerical methods in economics. The MIT press, 1998.
- [33] Justiniano, Alejandro, and Giorgio E. Primiceri. The time varying volatility of macroeconomic fluctuations. No. w12022. National Bureau of Economic Research, 2006.
- [34] Kim, Jinill, et al. "Calculating and using second-order accurate solutions of discrete time dynamic equilibrium models."
- [35] Kim, Jinill, and Sunghyun Henry Kim. "Spurious welfare reversals in international business cycle models." *Journal of International Economics* 60.2 (2003): 471-500. *Journal of Economic Dynamics and Control* 32.11 (2008): 3397-3414.
- [36] Kydland, Finn E., and Edward C. Prescott. "Time to build and aggregate fluctuations." *Econometrica: Journal of the Econometric Society* (1982): 1345-1370.
- [37] Lopes, J. S., and M. A. Beaumont. "ABC: a useful Bayesian tool for the analysis of population data." *Infection, Genetics and Evolution* 10.6 (2010): 825-832.

- [38] Marjoram, Paul, et al. "Markov chain Monte Carlo without likelihoods." *Proceedings of the National Academy of Sciences* 100.26 (2003): 15324-15328.
- [39] Nadaraya, Elizbar A. "On estimating regression." *Theory of Probability and Its Applications* 9.1 (1964): 141-142.
- [40] Pritchard, Jonathan K., Matthew Stephens, and Peter Donnelly. "Inference of population structure using multilocus genotype data." *Genetics* 155.2 (2000): 945-959.
- [41] Ruge-Murcia, Francisco. "Estimating nonlinear DSGE models by the simulated method of moments: With an application to business cycles." *Journal of Economic Dynamics and Control* 36.6 (2012): 914-938.
- [42] Schmitt-Grohe, Stephanie, and Martn Uribe. "Solving dynamic general equilibrium models using a second-order approximation to the policy function." *Journal of Economic Dynamics and Control* 28.4 (2004): 755-775.
- [43] Sims, Christopher. "Second order accurate solution of discrete time dynamic equilibrium models." Manuscript. Princeton: Princeton University (2000).
- [44] Sims, Christopher A. "Solving linear rational expectations models." *Computational Economics* 20.1 (2002): 1-20.
- [45] Sisson, S. A., Y. Fan, and Mark M. Tanaka. "Sequential monte carlo without likelihoods." *Proceedings of the National Academy of Sciences* 104.6 (2007): 1760-1765.
- [46] Smets, Frank, and Raf Wouters. "Comparing shocks and frictions in US and euro area business cycles: a Bayesian DSGE approach." *Journal of Applied Econometrics* 20.2 (2005): 161-183.
- [47] Smets, Frank, and Raf Wouters. "An estimated dynamic stochastic general equilibrium model of the euro area." *Journal of the European economic association* 1.5 (2003): 1123-1175.
- [48] Smets, Frank, and Raf Wouters. "Forecasting with a Bayesian DSGE model: an application to the euro area." *JCMS: Journal of Common Market Studies* 42.4 (2004): 841-867.
- [49] Watson, Geoffrey S. "Smooth regression analysis." *Sankhy: The Indian Journal of Statistics, Series A* (1964): 359-372.

Table 1: Prior distribution for the RBC parameters

Parameter	Distribution	1	2
β	<i>Beta</i>	0.95	0.02
γ	<i>Normal</i>	2	0.50
ρ_a	<i>Beta</i>	0.95	0.04
ρ_b	<i>Beta</i>	0.95	0.04
ρ_d	<i>Beta</i>	0.95	0.04
σ_a	<i>Gamma Inverse</i>	0.01	4
σ_b	<i>Gamma Inverse</i>	0.01	4
σ_d	<i>Gamma Inverse</i>	0.01	4

Prior distribution: Informative Prior

Table 2: RMSE, sample size=100 obs.

Methods	β	γ	ρ_a	ρ_b	ρ_d	σ_a	σ_b	σ_d
ABC-rej	0.01395	0.04079	0.01812	0.01566	0.01609	0.27268	0.22648	0.12772
ABC-ker	0.01456	0.04394	0.01871	0.01596	0.01666	0.27522	0.22939	0.13000
ABC-OLS	0.01406	0.06961	0.02131	0.02157	0.02014	0.27040	0.26608	0.16532
ABC-regr	0.01415	0.07220	0.02195	0.02180	0.02079	0.27223	0.26811	0.16567
ABC-HC	0.01920	0.10839	0.02755	0.03006	0.02448	0.26240	0.28406	0.22597
BLI	0.03729	0.05116	0.04154	0.03172	0.02695	0.67502	0.87365	0.30317

RMSE obtained in a Montecarlo experiment, 100 repetitions. The sample contains 100 observations. Case: High peristency and Informative Priors. ABC-rej= ABC-rejection, ABC-ker=ABC-rejection + kernel weighting, ABC-OLS= ABC + OLS Regression Step; ABC-regr= ABC-regression with Local Linear Regression, ABC-HC=ABC-regression + Correction for Heteroskedasticity

Table 3: RMSE, sample size=200 obs.

Methods	β	γ	ρ_a	ρ_b	ρ_d	σ_a	σ_b	σ_d
ABC-rej	0.01231	0.05162	0.01738	0.01691	0.01543	0.25187	0.22934	0.10386
ABC-ker	0.01332	0.05151	0.01798	0.01763	0.01606	0.25104	0.23001	0.10843
ABC-OLS	0.01237	0.06588	0.02006	0.02174	0.02197	0.24180	0.26175	0.13565
ABC-regr	0.01269	0.06876	0.01998	0.02186	0.02150	0.24271	0.26266	0.14553
ABC-HC	0.01655	0.09258	0.02385	0.02764	0.02650	0.22664	0.26542	0.20675
BLI	0.03418	0.10956	0.04294	0.05040	0.02682	0.58462	0.72849	0.59571

RMSE obtained in a Montecarlo experiment, 100 repetitions. The sample contains 200 observations. Case: High peristency and Informative Priors. ABC-rej= ABC-rejection, ABC-ker=ABC-rejection + kernel weighting, ABC-OLS= ABC + OLS Regression Step; ABC-regr= ABC-regression with Local Linear Regression, ABC-HC=ABC-regression + Correction for Heteroskedasticity

Table 4: RMSE, sample size=500 obs.

Methods	β	γ	ρ_a	ρ_b	ρ_d	σ_a	σ_b	σ_d
ABC-rej	0.01093	0.05065	0.02034	0.01764	0.01665	0.22934	0.27019	0.13820
ABC-ker	0.01110	0.05388	0.02042	0.01761	0.01669	0.22629	0.26472	0.14590
ABC-OLS	0.01081	0.08508	0.01765	0.01776	0.02078	0.22260	0.28040	0.19021
ABC-regr	0.01068	0.08764	0.01759	0.01755	0.02098	0.22184	0.28060	0.19879
ABC+HC	0.01205	0.11679	0.01875	0.01930	0.02520	0.20846	0.27512	0.26526
BLI	0.03742	0.06969	0.03863	0.03228	0.04938	0.58462	0.93056	0.53128

RMSE obtained in a Montecarlo experiment, 100 repetitions. The sample contains 500 observations. Case: High peristency and Informative Priors. ABC-rej= ABC-rejection, ABC-ker=ABC-rejection + kernel weighting, ABC-OLS= ABC + OLS Regression Step; ABC-regr= ABC-regression with Local Linear Regression, ABC-HC=ABC-regression + Correction for Heteroskedasticity

Table 5: OR100, sample size=100 obs.

Methods	β	γ	ρ_a	ρ_b	ρ_d	σ_a	σ_b	σ_d
ABC-rej	0.60627	0.81372	0.75159	0.80626	0.70472	0.44003	0.50961	0.70015
ABC-ker	0.67059	0.87735	0.76656	0.80266	0.80791	0.49469	0.58173	0.76644
ABC-OLS	0.36064	0.81120	0.72085	0.72093	0.75838	0.28645	0.39473	0.75102
ABC-regr	0.35319	0.80143	0.72284	0.72073	0.75071	0.28332	0.38944	0.74532
ABC-HC	0.55281	0.73430	0.66426	0.63165	0.68748	0.40299	0.42275	0.71252
BLI	0.04772	0.88188	0.66390	0.33132	0.34849	0.05231	-0.03619	0.43781

Overlapping Ratio obtained in a Montecarlo experiment, 100 repetitions. The sample contains 100 observations. Case: High peristency and Informative Priors. ABC-rej= ABC-rejection, ABC-ker=ABC-rejection + kernel weighting, ABC-OLS= ABC + OLS Regression Step; ABC-regr= ABC-regression with Local Linear Regression, ABC-HC=ABC-regression + Correction for Heteroskedasticity

Table 6: OR200, sample size=200 obs.

Methods	β	γ	ρ_a	ρ_b	ρ_d	σ_a	σ_b	σ_d
ABC-rej	0.57967	0.79299	0.73966	0.79252	0.67809	0.42241	0.47965	0.69985
ABC-ker	0.65300	0.87342	0.77590	0.78717	0.79739	0.48200	0.54320	0.76554
ABC-OLS	0.29644	0.82433	0.69746	0.68340	0.72241	0.22776	0.34818	0.74757
ABC-regr	0.29253	0.81414	0.69455	0.67834	0.72318	0.22574	0.34598	0.74664
ABC-HC	0.51729	0.77607	0.68972	0.67147	0.69955	0.39251	0.40124	0.71773
BLI	0.31990	0.77961	0.66926	0.62000	0.19545	0.12139	0.26556	0.43072

Overlapping Ratio obtained in a Montecarlo experiment, 100 repetitions. The sample contains 200 observations. Case: High peristency and Informative Priors. ABC-rej= ABC-rejection, ABC-ker=ABC-rejection + kernel weighting, ABC-OLS= ABC + OLS Regression Step; ABC-regr= ABC-regression with Local Linear Regression, ABC-HC=ABC-regression + Correction for Heteroskedasticity

Table 7: OR500, sample size=500 obs.

Methods	β	γ	ρ_a	ρ_b	ρ_d	σ_a	σ_b	σ_d
ABC-rej	0.50557	0.79718	0.68900	0.76053	0.64924	0.36181	0.44401	0.64670
ABC-ker	0.55337	0.86392	0.73684	0.76653	0.75746	0.40674	0.49992	0.73275
ABC-OLS	0.20349	0.79516	0.56611	0.58313	0.66112	0.13250	0.31733	0.69377
ABC-regr	0.20260	0.78619	0.56501	0.58256	0.65981	0.13104	0.31516	0.68630
ABC-HC	0.47321	0.76688	0.70070	0.74279	0.63753	0.34438	0.44529	0.67628
BLI	0.05993	0.83720	0.70892	0.31765	0.68671	0.11578	-0.04109	0.59651

Overlapping Ratio obtained in a Montecarlo experiment, 100 repetitions. The sample contains 500 observations. Case: High peristency and Informative Priors. ABC-rej= ABC-rejection, ABC-ker=ABC-rejection + kernel weighting, ABC-OLS= ABC + OLS Regression Step; ABC-regr= ABC-regression with Local Linear Regression, ABC-HC=ABC-regression + Correction for Heteroskedasticity

Par	Prior Distr	Prior Mean	Prior St.Dev.
β	<i>Beta</i>	0.997	0.001
θ	<i>Beta</i>	0.7	0.1
ϕ_y	<i>Gamma</i>	0.2	0.1
ϕ_π	<i>Gamma</i>	2.2	1
ρ_R	<i>Beta</i>	0.7	0.2
ϵ	<i>Gamma</i>	6	1
ϕ	<i>Gamma</i>	1	2
π	<i>Uniform</i>	1.002	1.007
$\bar{l}$	<i>Normal</i>	0	0.5
γ	<i>Normal</i>	0	0.5
ρ_A	<i>Beta</i>	0.70	0.20
ρ_G	<i>Beta</i>	0.70	0.20
ρ_U	<i>Beta</i>	0.80	0.10
σ_A	<i>InvGamma</i>	0.005	4
σ_G	<i>InvGamma</i>	0.005	4
σ_M	<i>InvGamma</i>	0.005	4
σ_U	<i>InvGamma</i>	0.005	4

Table 8: Prior distribution for the estimation of the newkeynesian model with the occasionally binding ZLB

Parameter	5% <i>CI</i>	Mean	95% <i>CI</i>
β	0.99591	0.996279	0.99671
θ	0.71878	0.764166	0.804385
ϕ_y	11 0.116811	0.158159	0.214062
ϕ_π	1.45885	1.85648	2.28268
ρ_R	0.643356	0.74771	0.836872
ϵ	6.74608	7.07615	7.45351
ϕ	0.15195	0.402306	0.759111
π	1.00207	1.00243	1.00295
ρ_A	0.602841	0.692637	0.775175
ρ_G	0.688226	0.735817	0.765022
ρ_U	0.825232	0.909938	0.978777
σ_A	0.00401074	0.0058359	0.00846274
σ_G	0.00394885	0.00479752	0.00609127
σ_M	0.00325742	0.00443026	0.00595078
σ_U	0.00401563	0.00548303	0.00745046
$\bar{l}$	0.0300085	0.144853	0.344083
γ	0.00592662	0.115779	0.312684

Table 9: Estimates for the Baseline sample (1966Q1-2014Q3), using only gaussian moments, ABC-SMC 10th iteration posterior mean and 5% and 95% credible interval values.

Parameter	5% <i>CI</i>	Mean	95% <i>CI</i>
β	0.99483	0.99520	0.99583
θ	0.66356	0.70538	0.73836
ϕ_y	0.10252	0.15220	0.19152
ϕ_π	1.49748	1.82866	2.25048
ρ_r	0.45842	0.51579	0.61003
ϵ	5.11345	5.56714	6.00983
ϕ	0.19322	0.59177	1.33045
π	1.00595	1.00621	1.00667
ρ_A	0.72697	0.82690	0.91660
ρ_G	0.67377	0.71112	0.76632
ρ_U	0.73527	0.83825	0.94086
σ_A	0.00325	0.00453	0.00576
σ_G	0.00337	0.00379	0.00423
σ_M	0.00338	0.00425	0.00494
σ_U	0.00352	0.00483	0.00623
l	0.00330	0.12036	0.29326
γ	0.00866	0.24216	0.53122

Table 10: Estimates for the Great Moderation sub-sample (1983Q1-2008Q3), using only gaussian moments, ABC-SMC 10th iteration posterior mean and 5% and 95% credible interval values.

Parameter	5% <i>CI</i>	Mean	95% <i>CI</i>
β	0.9967	0.9971	0.9974
θ	0.7403	0.7719	0.7973
ϕ_y	0.1385	0.1715	0.2073
ϕ_π	1.5818	1.817	2.2302
ρ_r	0.6532	0.7498	0.8106
ϵ	5.6922	5.9553	6.2826
ϕ	0.1547	0.5162	0.9885
π	1.0030	1.0035	1.0040
ρ_A	0.8421	0.9438	0.9923
ρ_G	0.7334	0.7806	0.8140
ρ_U	0.4984	0.5965	0.6743
σ_A	0.0030	0.0044	0.00634
σ_G	0.0038	0.0044	0.00508
σ_M	0.0039	0.00508	0.0067
σ_U	0.0031	0.0044	0.0055
l	0.0443	0.1680	0.3280
γ	0.1508	0.3175	0.5380

Table 11: Estimates for the Great Moderation +ZLB sub-sample (1983Q1-2014Q3), using only gaussian moments, ABC-SMC 10th iteration posterior mean and 5% and 95% credible interval values.

Parameter	5% <i>CI</i>	Mean	95% <i>CI</i>
β	0.99512	0.9953	0.99583
θ	0.69172	0.73243	0.8051
ϕ_y	0.11763	0.14271	0.1891
ϕ_π	1.28688	1.9525	2.5373
ρ_r	0.64005	0.7029	0.8352
ϵ	5.43814	5.8696	6.0763
ϕ	0.10567	0.3181	0.8044
π	1.0039	1.0046	1.00536;
ρ_A	0.77037	0.8747	0.9265
ρ_G	;0.70265	0.7492	0.8218
ρ_U	0.41432	0.50462	0.5395
σ_A	0.0037	0.0053	0.0073
σ_G	0.0046	0.0062	0.0070
σ_M	0.0035	0.0046	0.0056
σ_U	0.0043	0.00504	0.0068
l	0.0740	0.3899	0.5464
γ	0.01824	0.10519	0.40064

Table 12: Estimates for the Great Volatility II sub-sample (2001Q1-2014Q3), using only gaussian moments, ABC-SMC 10th iteration posterior mean and 5% and 95% credible interval values.

Parameter	5% <i>CI</i>	Mean	95% <i>CI</i>
β	0.99624	0.996852	0.997151
θ	0.706703	0.746625	0.791787
ϕ_y	0.114461	0.191332	0.264339
ϕ_π	1.49242	1.97145	2.36059
ρ_R	0.555455	0.6427	0.69883
ϵ	5.56305	5.88184	6.15413
ϕ	0.101903	0.353194	0.821647
π	1.00467	1.00515	1.00584
ρ_A	0.672021	0.748399	0.791278
ρ_G	0.719414	0.759741	0.789415
ρ_U	0.519773	0.59543	0.648594
σ_A	0.00337889	0.00437111	0.00567692
σ_G	0.00437548	0.00484554	0.00545708
σ_M	0.00336325	0.0043778	0.00548099
σ_U	0.0038067	0.0049121	0.00665101
l	0.0439899	0.202102	0.405959
γ	0.0383722	0.194152	0.348914

Table 13: Estimates for the Baseline sample (1966Q1-2014Q3), using gaussian and non-gaussian moments, ABC-SMC 10th iteration posterior mean and 5% and 95% credible interval values.

Parameter	5% <i>CI</i>	Mean	95% <i>CI</i>
β	0.99569	0.9962	0.9967
θ	0.7462	0.7864	0.8190
ϕ_y	0.17864	0.2308	0.2727
ϕ_π	2.2062	2.5943	2.90280
ρ_r	0.3969	0.4664	0.5124
ϵ	6.2492	6.6182	7.1278
ϕ	0.5348	1.6724	3.07298
π	1.0062	1.0064	1.00679
ρ_A	0.7309	0.8182	0.9135
ρ_G	0.7257	0.7617	0.8020
ρ_U	0.5355	0.6251	0.71412
σ_A	0.0039	0.0048	0.00587
σ_G	0.0034	0.0038	0.00441
σ_M	0.00314	0.0036	0.00428
σ_U	0.00404	0.0052	0.00680
l	0.5625	0.7337	0.9362
γ	0.0226	0.1072	0.2849

Table 14: Estimates for the Great Moderation sub-sample (1983Q1-2008Q3), using gaussian and non-gaussian moments, ABC-SMC 10th iteration posterior mean and 5% and 95% credible interval values.

Parameter	5% <i>CI</i>	Mean	95% <i>CI</i>
β	0.995834	0.996229	0.996659
θ	0.763125	0.794579	0.84167
ϕ_y	0.123546	0.178209	0.232961
ϕ_π	1.79105	2.20896	2.62851
ρ_r	0.606357	0.694911	0.790621
ϵ	5.07277	5.3655	5.68538
ϕ	0.152892	0.355819	0.806334
π	1.00274	1.00314	1.00371
ρ_A	0.678949	0.796617	0.861448
ρ_G	0.738915	0.761976	0.803161
ρ_U	0.728303	0.84731	0.919905
σ_A	0.00349434	0.00444531	0.00644926
σ_G	0.00387025	0.00477542	0.00580725
σ_M	0.00317167	0.00445699	0.00583731
σ_U	0.00435183	0.00615152	0.00815232
l	0.0148726	0.204401	0.416093
γ	0.00826553	0.176484	0.317272

Table 15: Estimates for the Great Moderation +ZLB sub-sample (1983Q1-2014Q3), using gaussian and non gaussian moments, ABC-SMC 10th iteration posterior mean and 5% and 95% credible interval values.

Parameter	5% <i>CI</i>	Mean	95% <i>CI</i>
β	0.9963	0.9968	0.9972
θ	0.6779	0.7133	0.7623
ϕ_y	0.1258	0.1706	0.2181
ϕ_π	1.3898	1.6264	2.0326
ρ_r	0.6158	0.7225	0.8099
ϵ	5.7602	6.1574	6.4772
ϕ	0.0867	0.2295	0.7533
π	1.0023	1.0028	1.0036
ρ_A	0.5577	0.6525	0.8684
ρ_G	0.7107	0.7650	0.7931
ρ_U	0.8613	0.9393	0.9914
σ_A	0.0042	0.0060	0.0076
σ_G	0.00280	0.0035	0.005
σ_M	0.0037	0.0046	0.0056
σ_U	0.0037	0.0048	0.0060
l	0.0139	0.1085	0.3500
γ	0.0292	0.1361	0.3746

Table 16: Estimates Great Volatility II sub-sample (2001Q1-2014Q3), using gaussian and non gaussian moments, ABC-SMC 10th iteration posterior mean and 5% and 95% credible interval values.

Figure 1: Distribution of the sample autocovariance for an AR(1) process with $\phi = 0.50$ for different sample sizes: from 50 to 2000 observations. The pink plane represents the population autocovariance.

Figure 2: Distribution of the sample autocovariance for an AR(1) process with $\phi = 0.99$ for different sample sizes: from 50 to 4000 observations. The pink plane represents the population autocovariance.

Figure 3: RMSE of the Montecarlo experiment: AR(1) process. Comparison among the ABC, HAC-BLI, Bootstrapping-BLI estimators, sample size=100. Different autocorrelations on the horizontal axis.

Figure 4: RMSE of the Montecarlo experiment: AR(1) process. Comparison among the ABC, HAC-BLI, Bootstrapping-BLI estimators, sample size=300. Different autocorrelations on the horizontal axis.

Figure 5: RMSE of the Montecarlo experiment: AR(1) process. Comparison among the ABC, HAC-BLI, Bootstrapping-BLI estimators, sample size=1000. Different autocorrelations on the horizontal axis.

Figure 6: Overlapping Ratios of the Montecarlo experiment: AR(1) process. Comparison among the ABC, HAC-BLI, Bootstrapping-BLI estimators, sample size=100. Different autocorrelations on the horizontal axis.

Figure 7: Overlapping Ratios of the Montecarlo experiment: AR(1) process. Comparison among the ABC, HAC-BLI, Bootstrapping-BLI estimators, sample size=300. Different autocorrelations on the horizontal axis.

Figure 8: Overlapping Ratios of the Montecarlo experiment: AR(1) process. Comparison among the ABC, HAC-BLI, Bootstrapping-BLI estimators, sample size=1000. Different autocorrelations on the horizontal axis.

Figure 9: Approximate posterior distributions obtained for the first 10 iterations of the ABC-SMC for the parameter ρ_U in an estimation exercise (Estimation of the subsample Great Moderation + ZLB, using gaussian moments).

Figure 10: 2-standard deviation Impulse responses of preference shock for 4 different sub-samples: Baseline (SW), Great Moderation (GM), Great Moderation+Zero Lower Bound (GM+ZLB), Great Volatility-II (GV). ABC-SMC is performed using only gaussian moments. Priors and 10-th iteration approximate posteriors are reported.

Figure 11: 2-standard deviation Impulse responses of preference shock for 4 different sub-samples: Baseline (SW), Great Moderation (GM), Great Moderation+Zero Lower Bound (GM+ZLB), Great Volatility-II (GV). ABC-SMC is performed using gaussian and non-gaussian moments. Priors and 10-th iteration approximate posteriors are reported.

Figure 12: Prior and posterior distributions for the newkeynesian model for 4 different sub-samples: Baseline (SW), Great Moderation (GM), Great Moderation+Zero Lower Bound (GM+ZLB), Great Volatility-II (GV). ABC-SMC is performed using only gaussian moments. Priors and 10-th iteration approximate posteriors are reported

Figure 13: Prior and posterior distributions for the newkeynesian model for 4 different sub-samples: Baseline (SW), Great Moderation (GM), Great Moderation+Zero Lower Bound (GM+ZLB), Great Volatility-II (GV). ABC-SMC is performed using only gaussian moments. Priors and 10-th iteration approximate posteriors are reported

Figure 14: Prior and posterior distributions for the newkeynesian model for 4 different sub-samples: Baseline (SW), Great Moderation (GM), Great Moderation+Zero Lower Bound (GM+ZLB), Great Volatility-II (GV). ABC-SMC is performed using only gaussian moments. Priors and 10-th iteration approximate posteriors are reported

Figure 15: Prior and posterior distributions for the newkeynesian model for 4 different sub-samples: Baseline (SW), Great Moderation (GM), Great Moderation+Zero Lower Bound (GM+ZLB), Great Volatility-II (GV). ABC-SMC is performed using gaussian and non-gaussian moments. Priors and 10-th iteration approximate posteriors are reported

Figure 16: Prior and posterior distributions for the newkeynesian model for 4 different sub-samples: Baseline (SW), Great Moderation (GM), Great Moderation+Zero Lower Bound (GM+ZLB), Great Volatility-II (GV). ABC-SMC is performed using gaussian and non-gaussian moments. Priors and 10-th iteration approximate posteriors are reported

Figure 17: Prior and posterior distributions for the newkeynesian model with for 4 different sub-samples: Baseline (SW), Great Moderation (GM), Great Moderation+Zero Lower Bound (GM+ZLB), Great Volatility-II (GV). ABC-SMC is performed using gaussian and non-gaussian moments. Priors and 10-th iteration approximate posteriors are reported